ECOLOGIA BALKANICA

2024, Vol. 16, Issue 1

June 2024

pp. 153-156

Short note

Summer diet of the red fox (Vulpes vulpes Linnaeus, 1785) in agricultural areas in South-Eastern Bulgaria

Alexander Petrov^{1*}, Ekaterina Pancheva²

¹University of Plovdiv "Paisii Hilendarski", Faculty of Biology, Department of Ecology and Environmental Conservation, Plovdiv, BULGARIA ²Trakia University, Agricultural Faculty, Stara Zagora, BULGARIA *Corresponding author: alex_petrov@uni-plovdiv.bg

Abstract. The summer diet of the red fox (*Vulpes vulpes* Linnaeus, 1785) in agricultural areas of South-Eastern Bulgaria was studied. Fecal samples were collected and processed in the laboratory of University of Plovdiv "Paisii Hilendarski". Insects, rodents, and fruits were the most commonly used dietary components. A relatively high percentage of reptiles were also found, which was observed in agricultural areas with a Mediterranean climate in Greece.

Key words: fecal samples, frequency of occurrence, dietary items.

Introduction

In the 21st century, many natural habitats have been modified by humans (Gosling, 2020). Much of the natural habitats are being converted into agricultural lands to meet the growing food needs of the human population (Blann, 2006). This forces many animals living in humaninfluenced areas to adapt to the new conditions.

The red fox (*Vulpes vulpes*) is among the preda-tors successfully surviving in such habitats, which manages to find food even in highly modified agricultural areas (Remonti et al., 2012). The high trophic plasticity of the predator allows it to adapt to the food resources of agroregions (Goldyn et al., 2003; Díaz-Ruiz et al., 2013).

Few studies have been conducted in Bulgaria on the food spectrum of this species in agricultural lands, and therefore this study aims to supplement the information on the summer diet of one of the most widespread predators in the country (Petrov et al., 2016; Petrov et al., 2024).

Materials and Methods

The survey was conducted from June 1, 2021 to August 31, 2021 in the area of Svilengrad municipality, South-Eastern Bulgaria (Fig. 1).

Agricultural production in the municipality is mainly aimed at growing tobacco, vineyards, cereals and fruits. Agriculture has a leading economic function in the municipality (Penin, 2007).

A total of 90 fecal samples (30 per month) were collected from areas with highly developed agriculture (the villages of Levka, Pastrogor, Lisovo, Mladinovo and Dipchevo). The samples were placed in plastic bags and processed in laboratory conditions at the University of Plovdiv "Paisii Hilendarski". Processing of the samples included deworming with 70% ethyl alcohol and water, washing through a sieve of 0.5 mm, determination of macrocomponents and their separation into 9 different groups.

The relative frequency of occurrence (RFO %) of all food components and food groups was calculated by dividing the number of occurrences of a certain food component by the sum of occurrences of all food components.



Fig. 1. Sample collection area.

Results and Discussion

Insects (32.32%), fruits (26.22%) and rodents (25.61%) were the most common food components found in the diet of the fox in South-Eastern Bulgaria, Svilengrad region during the summer season (Table 1). We agree with Goldyn et al. (2003) and Cancio et al. (2017) that the mosaic landscape, combining pastures, fields and fragments of oak forests, provides a variety of foraging opportunities. A relatively high percentage was also found in the captured reptiles (4.88%), which was also observed in other similar studies conducted in agricultural regions, but with a Mediterranean climate (Díaz-Ruiz et al., 2013; Bakaloudis et al., 2015).

Other studies of the fox diet in Bulgaria confirm that insects have a high frequency of

occurrence in the feces during the warm months, with only rodents and fruits being more common than them (Kyurkchiev, 2008; Petrov et al., 2016; Petrov et al., 2024).

Domestic birds (0.61%) and domestic mammals (2.44%) represented an extremely low percentage of the fox's diet during the study period. Much more preferred are rodents (25.61%) and mainly voles (14.2%). This shows the study predator's preference for hunting animals that humans find to be pests and its contribution to the regulation of rodent populations.

Seasonal fruits are also often used by the fox in this part of Bulgaria. The mulberry is one of the most frequently used food resources, which we explained by its accessibility and high distribution in the area. **Table 1.** Number of cases (n) and relative frequency of occurrence of food items in the faces (30 per month) of *Vulpes vulpes* in agricultural regions of South-Eastern Bulgaria.

Dietary items	June (n=30)		July (n=30)		August (n=30)		Total	
	S	RFO%	S	RFO%	S	RFO%	S	RFO%
Fruits	12	21.82	15	31.91	16	25.81	43	26.22
Grapes (Vitis sp.)	0	0.00	0	0.00	2	3.23	2	1.22
Mulberry (Morus sp.)	12	21.82	15	31.91	14	22.58	41	25.00
Domestic mammals	0	0.00	1	2.13	3	4.84	4	2.44
Goat (Capra domesticus)	0	0.00	0	0.00	1	1.61	1	0.61
Sheep (Ovis aries)	0	0.00	0	0.00	1	1.61	1	0.61
Wild mammals	0	0.00	1	2.13	1	1.61	2	1.22
Wild hare (Lepus europaeus)	0	0.00	0	0.00	1	1.61	1	0.61
Roe deer (Capreolus capreolus)	0	0.00	1	2.13	0	0.00	1	0.61
Rodents	14	25.45	10	21.28	18	29.03	42	25.61
Common wood mouse (Sylvaemus sylvaticus)	1	1.82	1	2.13	3	4.84	5	3.05
House mouse (Mus musculus)	3	5.45	2	4.26	5	8.06	10	6.10
Rat (Rattus norvegicus)	1	1.82	0	0.00	0	0.00	1	0.61
Vole (Microtus sp.)	7	12.73	7	14.89	9	14.52	23	14.02
Eadible dormouse (<i>Glis glis</i>)	2	3.64	0	0.00	1	1.61	3	1.83
Wild birds	1	1.82	0	0.00	1	1.61	2	1.22
Passeriformes	1	1.82	0	0.00	1	1.61	2	1.22
Domestic birds	0	0.00	0	0.00	1	1.61	1	0.61
Hen (Gallus gallus domesticus)	0	0.00	0	0.00	1	1.61	1	0.61
Reptiles	3	5.45	2	4.26	3	4.84	8	4.88
<i>Lacertilia</i> - undet.	3	5.45	2	4.26	3	4.84	8	4.88
Insects	20	36.36	16	34.04	17	27.42	53	32.32
Coleoptera	15	27.27	13	27.66	13	20.97	41	25.00
Orthoptera, Caelifera	3	5.45	3	6.38	4	6.45	10	6.10
Insecta - undet.	2	3.64	0	0.00	0	0.00	2	1.22
Other	5	9.09	2	4.26	2	3.23	9	5.49
Waste	3	5.45	1	2.13	1	1.61	5	3.05
Pebbles, grass	1	1.82	1	2.13	0	0.00	2	1.22
Egg shell	1	1.82	0	0.00	1	1.61	2	1.22
Total	55	100.00	47	100.00	62	100.00	164	100.00

References

Bakaloudis, D., Bontzorlos, V., Vlachos, C., Papakosta, M., Chatzinikos, E., Braziotis, S., & Kontsiotis, V. (2015). Factors affecting the diet of the red fox (*Vulpes vulpes*) in a heterogeneous Mediterranean landscape. *Turkish Journal of Zoology*, 39, 1151-1159. doi: 10.3906/zoo-1411-22

Blann, K. (2006). *Habitat in agricultural landscapes: How much is enough?*. Oregon. USA: West-Linn, 84 p.

Cancio, I., González-Robles, A., Bastida, J., Isla, J., Manzaneda, A., Salido, T., & Rey, P. (2017). Landscape degradation affects red fox (*Vulpes*)

vulpes) diet and its ecosystem services in the threatened *Ziziphus lotus* scrubland habitats of semiarid Spain. *Journal of Arid Environments*, 145, 24–34. doi: 10.1016/j.jaridenv.2017.05.004

Díaz-Ruiz, F., Delibes-Mateos, M., García-Moreno, J., López-Martín, J., Ferreira, C., & Ferreras, P. (2013). Biogeographical patterns in the diet of an opportunistic predator: the red fox *Vulpes vulpes* in the Iberian Peninsula. *Mammal Review*, 43, 59–70. doi: 10.1111/j.1365-2907.2011.00206.x

Goldyn, B., Hromada, M., Surmacki, A., & Tryjanowski, P. (2003). Habitat use and diet of the red fox *Vulpes vulpes* in an agricultural landscape in Poland. Zeitschrift fur jagdwissenschaft, 49, 1-10. doi: 10.1007/BF02189737

Gosling, J., Jones, M., Arnell, A., Watson, J., Venter, O., Baquero, A., & Burgess, N. (2020). A global mapping template for natural and modified habitat across terrestrial Earth. *Biological Conservation*, 250, 108-674.

doi: 10.1016/j.biocon.2020.108674

Kidawa, D., & Kowalczyk, R. (2011). The effects of sex, age, season and habitat on diet of the red fox *Vulpes vulpes* in northeastern Poland. *Acta Theriologica*, 56, 209–218. doi: 10.1007/s13364-011-0031-3

Kyurkchiev, S. (2008). Godishen hranitelen spektar na chervenata lisitsa (*Vulpes vulpes* L.) v Tzentralni Rhodopi. [Annual food spectrum of the Red Fox (*Vulpes vulpes* L.) in Central Rhodopi]. Proceedings of the Jubilee scientific conference on Ecology, Plovdiv, pp. 200-207. (In Bulgarian).

Penin, R. (2007). *Natural geography of Bulgaria*. Sofia, Bulgaria: Bulvest 2000, p. 280. (In Bulgarian) Petrov, P., Popova, E., & Zlatanova, D. (2016). Niche partitioning among the Red Fox *Vulpes vulpes* (L.), Stone Marten *Martes foina* (Erxleben) and Pine marten *Martes martes* (L.) in Two Mountains in Bulgaria. *Acta zoologica bulgarica*, 68(3), 375-390.

Petrov, A., Peeva, S., & Mollov, I. (2024) What's for dinner? Diet and trophic niche overlap in two sympatric carnivores in agricultural and nearnatural habitats in Bulgaria. *Ecologia Balkanica*, 16, 21-29.

Remonti, L., Balestrieri, A., Ruiz-González, A., Gómez-Moliner, B., Capelli, E., & Prigioni, C. (2012). Intraguild dietary overlap and its possible relationship to the coexistence of mesocarnivores in intensive agricultural habitats. *Population Ecology*, 54, 521-532. doi: 10.1007/S10144-012-0337-2

> Received: 13.05.2023 Accepted: 09.06.2024